

Blood Sugar Blaster Research:

Vitamin C

- Afkhami-Ardekani M, Shojaoddiny-Ardekani A. Effect of vitamin C on blood glucose, serum lipids & serum insulin in type 2 diabetes patients. *Indian J Med Res.* 2007;126(5):471-4.
- Dakhale GN1, Chaudhari HV, Shrivastava M. Supplementation of vitamin C reduces blood glucose and improves glycosylated hemoglobin in type 2 diabetes mellitus: a randomized, double-blind study. *Adv Pharmacol Sci.* 2011;2011:195271.
- M. A. Ardekani and A. S. Ardekani, "Effect of vitamin C on blood glucose, serum lipids & serum insulin in type II diabetes patients," *Indian Journal of Medical Research*, vol. 126, no. 5, pp. 471–474, 2007
- N. Bishop, C. J. Schorah, and J. K. Wales, "The effect of vitamin C supplementation on diabetic hyperlipidaemia: a double blind, crossover study," *Diabetic Medicine*, vol. 2, no. 2, pp. 121–124, 1995.
- P. Sridulyakul, D. Chakraphan, and S. Patumraj, "Vitamin C supplementation could reverse diabetes-induced endothelial cell dysfunction in mesenteric microcirculation in STZrats," *Clinical Hemorheology and Microcirculation*, vol. 34, no. 1-2, pp. 315–321, 2006.
- Ashor AW, Werner AD, Lara J. Effects of vitamin C supplementation on glycaemic control: a systematic review and meta-analysis of randomised controlled trials. *Eur J Clin Nutr.* 2017 Mar 15.
- S. J. Davie, B. J. Gould, and J. S. Yudkin, "Effect of vitamin C on glycosylation of proteins," *Diabetes*, vol. 41, no. 2, pp. 167–173, 1992.

Vitamin E

- Suksomboon N, Poolsup N, Sinprasert S. Effects of vitamin E supplementation on glycaemic control in type 2 diabetes: systematic review of randomized controlled trials. *Vit J Clin Pharm Ther.* 2011 Feb;36(1):53-63. [Article](#).
- Goldenstein H, Levy NS, Lipener YT, Levy AP. Patient selection and vitamin E treatment in diabetes mellitus. *Expert Rev Cardiovasc Ther.* 2013 Mar;11(3):319-26. [Article](#).
- de Oliveira A. M., Rondó P. H., Luzia L. A. The effects of lipoic acid and α -tocopherol supplementation on the lipid profile and insulin sensitivity of patients with type 2 diabetes mellitus: a randomized, double-blind, placebo-controlled trial. *Diabetes Res Clin Pract.* 2011 May;92(2):253-60.
- Park S, Choi SB (2002) Effects of alpha-tocopherol supplementation and continuous subcutaneous insulin infusion on oxidative stress in Korean patients with type 2 diabetes. *Am J Clin Nutr* 75: 728–733
- Ble-Castillo JL, Carmona-Diaz E, Mendez JD, Larios-Medina FJ, Medina-Santillan R, et al. (2005) Effect of alpha-tocopherol on the metabolic control and oxidative stress in female type 2 diabetics. *Biomed Pharmacother* 59: 290–295
- Boshtam M, Rafiei M, Golshadi ID, Ani M, Shirani Z, et al. (2005) Long term effects of oral vitamin E supplement in type II diabetic patients. *Int J Vitam Nutr Res* 75: 341–346
- Ceriello A, Giugliano D, Quatraro A, Donzella C, Dipalo G, et al. (1991) Vitamin E reduction of protein glycosylation in diabetes. New prospect for prevention of diabetic complications? *Diabetes Care* 14: 68–72
- Liu S, Lee IM, Song Y, Van Denburgh M, Cook NR, et al. (2006) Vitamin E and risk of type 2 diabetes in the women's health study randomized controlled trial. *Diabetes* 55: 2856–2862
- Suksomboon N, Poolsup N, Sinprasert S (2011) Effects of vitamin E supplementation on glycaemic control in type 2 diabetes: systematic review of randomized controlled trials. *J Clin Pharm Ther* 36: 53–63
- Xu R, Zhang S, Tao A. Influence of vitamin E supplementation on glycaemic control: a metaanalysis

of randomized controlled trials. *PLoS One*. 2014 Apr 16;9(4):e95008. [Article](#).
Paolisso G, D'Amore A, Galzerano D, Balbi V, Giugliano D, et al. (1993) Daily vitamin E supplements improve metabolic control but not insulin secretion in elderly type II diabetic patients. *Diabetes Care* 16: 1433–1437
Gazis A, White DJ, Page SR, Cockcroft JR (1999) Effect of oral vitamin E (alpha-tocopherol) supplementation on vascular endothelial function in Type 2 diabetes mellitus. *Diabet Med* 16: 304–311

Biotin

Albarracín CA, Fuqua BC, Evans JL, Goldfine ID. Chromium picolinate and biotin combination improves glucose metabolism in treated, uncontrolled overweight to obese patients with type 2 diabetes. *Diabetes Metab Res Rev*. 2008 Jan-Feb;24(1):41-51.
McCarty MF. Supplementation with Phycocyanobilin, Citrulline, Taurine, and Supranutritional Doses of Folic Acid and Biotin-Potential for Preventing or Slowing the Progression of Diabetic Complications. *Healthcare (Basel)*. 2017 Mar 14;5(1).
McCarty MF. In type 1 diabetics, high-dose biotin may compensate for low hepatic insulin exposure, promoting a more normal expression of glycolytic and gluconeogenic enzymes and thereby aiding glycemic control. *Med Hypotheses*. 2016 Oct;95:45-48.
Hemmati M, Babaei H, Abdolsalehei M. Survey of the effect of biotin on glycemic control and plasma lipid concentrations in type 1 diabetic patients in kermanshah in iran (2008-2009). *Oman Med J*. 2013 May;28(3):195-8.
The effect of chromium picolinate and biotin supplementation on glycemic control in poorly controlled patients with type 2 diabetes mellitus: a pla... - PubMed – NCBI

Magnesium

J. Eriksson and A. Kohvakka, “Magnesium and ascorbic acid supplementation in diabetes mellitus,” *Annals of Nutrition and Metabolism*, vol. 39, no. 4, pp. 217–223, 1995.
White JR Jr, Campbell RK. Magnesium and diabetes: a review. *Ann Pharmacother*. 1993 Jun; 27(6):775-80. [Article](#).
Medicine (Baltimore). 2017 Mar;96(11):e6352. doi: 10.1097/MD.0000000000006352.
Shahbah D1, Hassan T, Morsy S. Oral magnesium supplementation improves glycemic control and lipid profile in children with type 1 diabetes and hypomagnesaemia. *Medicine (Baltimore)*. 2017 Mar;96(11):e6352.
Rodríguez-Morán M, Guerrero-Romero F. Oral magnesium supplementation improves insulin sensitivity and metabolic control in type 2 diabetic subjects: a randomized double-blind controlled trial. *Diabetes Care*2003;26:1147–52.
Simental-Mendía LE, Sahebkar A, Rodríguez-Morán M, et al. A systematic review and metaanalysis of randomized controlled trials on the effects of magnesium supplementation on insulin sensitivity and glucose control. *Pharmacol Res* 2016;111:272–82.

Zinc

Effects of zinc supplementation on diabetes mellitus: a systematic review and meta-analysis. Jayawardena R, Ranasinghe P, Galappathy P, Malkanthi R, Constantine G, Katulanda P. *Diabetol Metab Syndr*. 2012 Apr 19;4(1):13.
Shidfar F, Aghasi M, Vafa M, Heydari I, Hosseini S, Shidfar S. Effects of combination of zinc and vitamin A supplementation on serum fasting blood sugar, insulin, apoprotein B and apoprotein AI in patients with type i diabetes. *International Journal of Food Sciences and Nutrition*. 2010;61:182–191.
Al-Maroofof RA, Al-Sharbatti SS. Serum zinc levels in diabetic patients and effect of zinc supplementation on glycemic control of type 2 diabetics. *Saudi Medical Journal*. 2006;27:344–350.

de Sena KC, Arrais RF, das Gracas Almeida M, de Araujo DM, dos Santos MM, de Lima VT, de Fatima Campos Pedrosa L: Effects of zinc supplementation in patients with type 1 diabetes. *Biol Trace Elem Res.* 2005;105:1–9.

Anderson RA, Roussel AM, Zouari N, Mahjoub S, Matheau JM, Kerkeni A. Potential antioxidant effects of zinc and chromium supplementation in people with type 2 diabetes mellitus. *J Am Coll Nutr.* 2001;20:212–218.

Gunasekara P, Hettiarachchi M, Liyanage C, Lekamwasam S. Effects of zinc and multimineral vitamin supplementation on glycemic and lipid control in adult diabetes. *Diabetes, Metabolic Syndrome and Obesity: Targets and Therapy.* 2011;4:53–60.

Manganese

Koh ES, Kim SJ, Yoon HE. Association of blood manganese level with diabetes and renal dysfunction: a cross-sectional study of the Korean general population. *BMC Endocr Disord.* 2014 Mar 8;14:24. [Article](#).

Oral Magnesium Supplementation Improves Insulin Sensitivity and Metabolic Control in Type 2 Diabetic Subjects. [[ADA](#)]

Chromium

[No authors listed]. A scientific review: the role of chromium in insulin resistance. *Diabetes Educ.* 2004;Suppl:2-14.

Paiva AN, Lima JG, Medeiros AC. Beneficial effects of oral chromium picolinate supplementation on glycemic control in patients with type 2 diabetes: A randomized clinical study. *J Trace Elem Med Biol.* 2015 Oct;32:66-72. [Article](#).

Effects of chromium supplementation on fasting insulin levels and lipid parameters in healthy, non-obese young subjects. [[PubMed](#)]

Albarracin CA, Fuqua BC, Evans JL, Goldfine ID. Chromium picolinate and biotin combination improves glucose metabolism in treated, uncontrolled overweight to obese patients with type 2 diabetes. *Diabetes Metab Res Rev.* 2008 Jan-Feb;24(1):41-51.

Singer GM, Geohas J. The effect of chromium picolinate and bio<n supplementa<on on glycemic control

in poorly controlled pa<ents with type 2 diabetes mellitus: a placebo-controlled, double-blinded, randomized trial. *Diabetes Technol Ther* 2006. Dec;8(6):636-643

Anderson RA, Roussel AM, Zouari N, Mahjoub S, Matheau JM, Kerkeni A. Poten<al an<oxidant effects of

zinc and chromium supplementa<on in people with type 2 diabetes mellitus. *J Am Coll Nutr.* 2001;20:212–218.

Balk EM, Tatsioni A, Lichtenstein AH. Effect of chromium supplementation on glucose metabolism and lipids: a systematic review of randomized controlled trials. *Diabetes Care.* 2007 Aug;30(8):2154-63.

Wang X, Zhang M, Lui G. Associations of Serum Manganese Levels with Prediabetes and Diabetes among ≥60-Year-Old Chinese Adults: A Population-Based Cross-Sectional Analysis. *Nutrients.* 2016 Aug 13;8(8). [Article](#).

Banaba Leaf extract

Miura T, Takagi S, Ishida T. Management of Diabetes and Its Complications with Banaba (*Lagerstroemia speciosa* L.) and Corosolic Acid. *Evid Based Complement Alternat Med.* 2012;2012:871495.

Management of Diabetes and Its Complications with Banaba (*Lagerstroemia speciosa* L.) and Corosolic Acid. [[PubMed](#)] [[PubMed](#)]

Guggul

Sarup P1, Bala S1, Kamboj S1. Pharmacology and Phytochemistry of Oleo-Gum Resin of *Commiphora wightii* (Guggulu). *Scientifica* (Cairo). 2015;2015:138039. [Article](#).
R. Bellamkonda, K. Rasineni, S. R. Singareddy et al., “Antihyperglycemic and antioxidant activities of alcoholic extract of *Commiphora mukul* gum resin in streptozotocin induced diabetic rats,” *Pathophysiology*, vol. 18, no. 4, pp. 255–261, 2011.

Bitter Melon

Alam MA, Uddin R, Subhan N. Beneficial Role of Bitter Melon Supplementation in Obesity and Related Complications in Metabolic Syndrome. *J Lipids*. 2015; 2015: 496169.
Krawinkel M. B., Keding G. B. Bitter gourd (*Momordica charantia*): a dietary approach to hyperglycemia. *Nutrition Reviews*. 2006;64(7):331–337.
Tsai C.-H., Chen E. C.-F., Tsay H.-S., Huang C.-J. Wild bitter gourd improves metabolic syndrome: a preliminary dietary supplementation trial. *Nutrition Journal*. 2012;11
Fuangchan A., Sonthisombat P., Seubnukarn T., et al. Hypoglycemic effect of bitter melon compared with metformin in newly diagnosed type 2 diabetes patients. *Journal of Ethnopharmacology*. 2011;134(2):422–428.
Leung L., Birtwhistle R., Kotecha J., Hannah S., Cuthbertson S. Anti-diabetic and hypoglycaemic effects of *Momordica charantia* (bitter melon): a mini review. *British Journal of Nutrition*. 2009;102(12):1703–1708.

Licorice root extract

Cheng HS, Kong JM, Ng AX. Novel Inhibitory Effects of Glycyrrhizic Acid on the Accumulation of Advanced Glycation End Product and Its Receptor Expression. *Nat Prod Bioprospect*. 2014 Dec;4(6):325-33.
Kuroda M, Mimaki Y, Sashida Y. Phenolics with PPAR-gamma ligand-binding activity obtained from licorice (*Glycyrrhiza uralensis* roots) and ameliorative effects of glycyrrin on genetically diabetic KK-A(y) mice. *Bioorg Med Chem Lett*. 2003 Dec 15;13(24):4267-72. [Article](#).
Chia YY, Liong SY, Ton SH. Amelioration of glucose homeostasis by glycyrrhizic acid through gluconeogenesis rate-limiting enzymes. *Eur J Pharmacol*. 2012 Feb 29;677(1-3):197-202.

Cinnamon Bark Powder

Mang B, Wolters M, Schmitt B, Kelb K, Lichtinghagen R, Stichtenoth DO, et al. Effects of a cinnamon extract on plasma glucose, HbA1c, and serum lipids in diabetes mellitus type 2. *Eur J Clin Invest*. 2006;36:340–4.
Crawford P. Effectiveness of cinnamon for lowering hemoglobin A1C in patients with type 2 diabetes: a randomized, controlled trial. *J Am Board Fam Med*. 2009;22:507–12. [[PubMed](#)]
Akilen R, Tsiami A, Devendra D, Robinson N. Glycated haemoglobin and blood pressure lowering effect of cinnamon in multi-ethnic Type 2 diabetic patients in the UK: a randomized, placebo-controlled, double-blind clinical trial. *Diabet Med*. 2010;27:1159–67.
Itschuler JA, Casella SJ, MacKenzie TA, Curtis KM. The effect of cinnamon on A1C among adolescents with type 1 diabetes. *Diabetes Care*. 2007;30:813–6. [[PubMed](#)]
Blevins SM, Leyva MJ, Brown J, Wright J, Scofield RH, Aston CE. Effect of cinnamon on glucose and lipid levels in non insulin-dependent type 2 diabetes. *Diabetes Care*. 2007;30:2236–7. [[PubMed](#)]
Mettler S, Schwarz I, Colombani PC. Additive postprandial blood glucose-attenuating and satiety-enhancing effect of cinnamon and acetic acid. *Nutr Res*. 2009;29:723–7. [[PubMed](#)]
Cinnamon: potential role in the prevention of insulin resistance, metabolic syndrome, and type 2 diabetes.– [[PubMed](#)]
Cinnamon improves glucose and lipids of people with type 2 diabetes.– [[PubMed](#)]

Solomon TP, Blannin AK. Changes in glucose tolerance and insulin sensitivity following 2 weeks of daily cinnamon ingestion in healthy humans. *Eur J Appl Physiol.* 2009;105:969– [PubMed]

Gymnema Sylvestre Leaf Powder

Kanetkar P, Singhal R, Kamat M. *Gymnema sylvestre*: A Memoir. *J Clin Biochem Nutr.* 2007 Sep; 41(2): 77–81.

Persaud S.J., Al-Majed H., Raman A., Jones P.M. *Gymnema sylvestre* stimulates insulin release in vitro by increased membrane permeability. *J. Endocrinol.* 1999;163:207–212.

Khare A.K., Tondon R.N., Tewari J.P. Hypoglycemic activity of an indigenous drug *Gymnema sylvestre* in normal and diabetic persons. *Ind. J. Physiol. Pharmacol.* 1983;27:257–261.

Antidiabetic effect of a leaf extract from *Gymnema sylvestre* in non-insulin-dependent diabetes mellitus patients. - PubMed - NCBI." National Center for Biotechnology Information. U.S.

National Library of Medicine, n.d. Web. 18 June 2017.

Yarrow Flowers Powder

Yazdanparast R, Ardestani A, Jamshidi S. Experimental diabetes treated with *Achillea santolina*: effect on pancreatic oxidative parameters. *J Ethnopharmacol.* 2007 May 30;112(1):13-8. [Article](#).

Tiwari P, Mishra BN, Sangwan NS. Phytochemical and pharmacological properties of *Gymnema sylvestre*: an important medicinal plant. *Biomed Res Int.* 2014;2014:830285. [Article](#).

Yazdanparast R, Ardestani A, Jamshidi S. Experimental diabetes treated with *Achillea santolina*: effect on pancreatic oxidative parameters. *J Ethnopharmacol.* 2007 May 30;112(1):13-8.

Shanmugasundaram ERB, Rajeswari G, Baskaran K, Kumar BRR, Shanmugasundaram KR, Ahmath BK. Use of *Gymnema sylvestre* leaf extract in the control of blood glucose in insulin independent

diabetes mellitus. *Journal of Ethnopharmacology.* 1990;30(3):281–294.

Baskaran K, Ahmath BK, Shanmugasundaram KR, Shanmugasundaram ERB. Antidiabetic effect of a leaf extract from *Gymnema sylvestre* in non-insulin-dependent diabetes mellitus patients. *Journal of Ethnopharmacology.* 1990;30(3):295–305.

Cayenne pepper Powder

Khan I, Ahmad H, Ahmad B. Anti-glycation and anti-oxidation properties of *Capsicum frutescens* and *Curcuma longa* fruits: possible role in prevention of diabetic complication. *Pak J Pharm Sci.* 2014 Sep;27(5):1359-62. [Article](#).

Chaiyasit K, Khovidhunkit W, Wittayalertpanya S. Pharmacokinetic and the effect of capsaicin in *Capsicum frutescens* on decreasing plasma glucose level. *J Med Assoc Thai.* 2009 Jan;92(1): 108-13. [Article](#).

Tolan I, Ragoobirsingh D, Morrison EY. Isolation and purification of the hypoglycaemic principle present in *Capsicum frutescens*. *Phytother Res.* 2004 Jan;18(1):95-6.

Juniper berry powder

Sánchez de Medina F, Gámez MJ, Jiménez I, Jiménez J, Osuna JI, Zarzuelo A. Hypoglycemic activity of juniper “berries” *Planta Med.* 1994;60:197–200.

White mulberry leaf powder

Kimura T., Nakagawa K., Kubota H., Kojima Y., Yamagishi K., Oita S., Oikawa S., Miyazawa T. Food-grade mulberry powder enriched with 1-deoxynojirimycin suppresses the elevation of postprandial blood glucose in humans. *J. Agric. Food Chem.* 2007;55:5869–5874.

Mulberry-extract improves glucose tolerance and decreases insulin concentrations in normoglycaemic adults: Results of a randomised double-blind placebo-controlled study.

Lown M, Fuller R, Lightowler H, Fraser A, Gallagher A, Stuart B, Byrne C, Lewith G. PLoS One. 2017 Feb 22;12(2). [Article](#).

Mulberry leaf extract improves postprandial glucose response in prediabetic subjects: a randomized, double-blind placebo-controlled trial.

Kim JY, Ok HM, Kim J, Park SW, Kwon SW, Kwon O.

J Med Food. 2015 Mar;18(3):306-13. [Article](#).

Effect of mulberry (*Morus indica* L.) therapy on plasma and erythrocyte membrane lipids in patients with type 2 diabetes. [PubMed](#)

Influence of Mulberry Leaf Extract on the Blood Glucose and Breath Hydrogen Response to Ingestion of 75 g Sucrose by Type 2 Diabetic and Control Subjects. [ADA](#)

Vanadium

Badmaev V, Prakash S, Majeed M. Vanadium: a review of its potential role in the fight against diabetes. *J Altern Complement Med*. 1999 Jun;5(3):273-91.

Jacques-Camarena O, González-Ortiz M, Martínez-Abundis E, López-Madrueño JF, Medina-Santillán R. Effect of vanadium on insulin sensitivity in patients with impaired glucose tolerance. *Ann Nutr Metab*. 2008;53(3-4):195-8. [Article](#).

Thompson KH, Lichter J, LeBel C. Vanadium treatment of type 2 diabetes: a view to the future. *J Inorg Biochem*. 2009 Apr;103(4):554-8. [Article](#).

Thompson KH, Orvig C. Vanadium in diabetes: 100 years from Phase 0 to Phase I. *J Inorg Biochem*. 2006 Dec;100(12):1925-35. [Article](#).

Alpha lipoic acid

Saeid Golbidi, Mohammad Badran, and Ismail Laher. Diabetes and Alpha Lipoic Acid. *Front Pharmacol*. 2011; 2: 69.

Ansar H., Mazloom Z., Kazemi F., Hejazi N. (2011). Effect of alpha-lipoic acid on blood glucose, insulin resistance and glutathione peroxidase of type 2 diabetic patients. *Saudi Med. J.* 32, 584–588.

de Oliveira A. M., Rondó P. H., Luzia L. A., D'Abronzio F. H., Illison V. K. (2011). The effects of lipoic acid and α -tocopherol supplementation on the lipid profile and insulin sensitivity of patients with type 2 diabetes mellitus: a randomized, double-blind, placebo-controlled trial. *Diabetes Res. Clin. Pract.* 92, 253–26010.

Kamenova P. (2006). Improvement of insulin sensitivity in patients with type 2 diabetes mellitus after oral administration of alpha-lipoic acid. *Hormones (Athens)* 5, 251–258

L-taurine

Franconi F, Loizzo A, Ghirlanda G, Seghieri G. Taurine supplementation and diabetes mellitus. *Curr Opin Clin Nutr Metab Care*. 2006;9:32–36.

Ito T, Schaffer SW, Azuma J. The potential usefulness of taurine on diabetes mellitus and its complications. *Amino Acids*. 2012 May; 42(5): 1529–1539.

Chauncey KB, Tenner TE, Jr, Lombardini JB, Jones BG, Brooks ML, Warner RD, Davis RL, Ragain RM. The effect of taurine supplementation on patients with type 2 diabetes mellitus. *Adv Exp Med Biol*. 2003;526:91–96.

Elizarova EP, Nedosugova LV. First experiments in taurine administration for diabetes mellitus. The effect on erythrocyte membranes. *Adv Exp Med Biol*. 1996;403:583–588.